REFERENCES

1. Pandit JK, Targeted Devices for Periodontal disease, Chapter-6 in controlled and novel drug delivery, Jain MK. First edition, C.B.S. Publishers and Distribution, New Delhi, 1997, 130-146.
2. Eickholz P,Dannewitz B, Ti-Sun Kim, Antibiotics in periodontal therapy. Clinical and research reports. 2005, 2(4):235-251.
3. Poveda Roda R , Vicente Bagan J, Maria J , Carbonell Pastor E. Antibiotic use in dental practice. A review, Med Oral Patol Oral Cir Bucal 2007;12,186.
4. Aulton ME, Pharmaceutics, the Science of Dosage Forms Design. 2nd ed. Harcourt Publishers Limited; 2002. 88.
5. Subhas G, Local drug delivery in periodontics, Indian Journal of Dental Sciences. Jan 2010,2(1).
6. Swati R, Sandeep W, Swaroop L, *In Situ* Gel Formulation of Ornidazole for the Treatment of Periodontal Disease, Current Pharma Research, 2010,1(1).
7. Ananta c, Sujoy D, Satish D, Sumit K, Abhishak K, Development and Characterization Buccoadhesive Film of Ciprofloxacin Hydrochloride, International Journal of PharmTech Research, 2010,2(2),1050-1057.
8. Jongjan M, Juree C, Thawatchai P, Zinc Oxide Gels for Periodontitis Treatment, Journal of Metals, Materials and Minerals, 2010.20(3),159-163.
9. Bansal K, Rawat M, Jain A, Rajput A, Chaturvedi T , Singh S, Development of starnidazole mucoadhesive gel for treatment of periodontitis, AAPS Pharm SciTech, 2009. 10(3).
10. Pragati S, Ashok S., Kuldeep S. Recent advances in periodontal drug delivery systems, International Journal of Drug Delivery, 2009 1-14.
11. Mastiholimath VS, Dandagi PM, Gadad AP, Patil MB, Manvi FV, and Chandur VK, Formulation and evaluation of ornidazole dental implants for periodontitis. Ind. J. Pharm. Sci. 2006; 68.

12. Alka Ahuja, Ali J, Shareef A, and Khar RK, Formulation and development of targeted retentive device of the treatment of periodontal infection with amoxycillin trihydrate. Ind. J. Pharm. Sci. 2006, 442.

13. Giuseppe and perinetti, clinical and microbiological effects of subgingivalasministration of two active gels on persistant pocket of chronic periodontitis patients , journal of clinical periodontology, 2004, 31(4),282.

14. Perugini P, Genta I, Conti B, Modena T,pavanetto F, periodontal deleviry of ipriflavone: new chitosan/PLGA film deliverysystem for a lipophilic drug, International journal of pharmaceutics.2003, 252(1), 1-9.

15. Nagaraju R, Udupa N, Verma BRR, Tinidazole concentration in Human Gingival Crevicular Fluid after insertion of Biodegradable Dental Implants, Indian Journal of Pharma Sci, 2003, 540.

16. Sharma N, and Galustains H, antiplaque and antigingivitis effectiveness of hexidine mouthwash, journal of clinical periodontology, 2003,30; 590-594.

17. Behnen, Micheal J, Antibacterial activity of several calcium hydroxides preparation in root canal dentine, J. Endodont, 2001,27(12), 765.

18. Karnik a, vania p, Novel gel-sol injection of Doxacyclin hydrochloride for the treatment of periodontitis.scientific poster presentation 53rd IPC 2001.

19. Kargill B and Kadir T, the antibacterial effect of ornidazole on primary molars with infected pulps, experimental chemotherapy, 2001, 47 ,203-207.

20. Vyas S, Shiorkar V,Mishra V, controlled and targeted drug delivery strategies toward intraperiodontalpocket diseases, journal of clinical pharma therapy pharmacy and therapeutics, 2000,25,21.

21. Reinhardt R, Maze G, Poynl J, and Weber D, Gingival crevicular fluid tetracycline release from biodegradable gels. J. Dent. Res., 1995, 74, 531.

22. Giordano J, and Loasche WJ, Effect of local delivery of antimicrobial agent to tooth refractory to debridement plus systemic antimicrobials. J. Dent. Res., 1995, 26.

23. Walker C, Magnusson I, And Heft A, Effect of metronidazole containing gel on the periodontal microflora. J. Dent. Res., 1995, 74, 481.

24. Polson AM, Garrett S, Stoller N, and Nandit C, Multicentre study of sub-gingival doxycyclin in treatment of periodontitis J. Dent. Res., 1995, 74, 26.

25. Lehmann B, Mombelli A, Tonetti M, and Lang NP, Clinical response to local delivery of tetracycline in relation to overall conditions of other teeth. J. Dent. Res., 1995, 74, 481.

26. Killoy WJ, Rapely JW, Drisko CL, Yonke ML, and Ridenhoujr LS, Twelve month comparison of tetracycline fiber treatment vs periodontal maintenance. J. Dent. Res., 1995, 74, 480.

27. Drisko CL, and Charles M, Evaluation of periodontal treatments using controlled release tetracycline fibers- clinical response. J. Periodontal, 1995, 66, 692.

28. Sabuski D, Asthon P, Abrams H, and Kupp L, Evaluation of two types of biodegradable systems for local delivery. J. Dent. Res., 1994, 73, 308.

29. Jones AA, Korman KS, Newbold DA, and Manwell MA, Clinical and microbiological effects of controlled release locally delivered minocycline in periodontitis. J. Periodontal., 1994, 65, 1058.

30. Lowenguth RM, and Gandini L, Effects of sub-gingival delivery of tetracycline on the microflora in localized juvenile periodontitis. J. Dent. Res., 1993, 72, 160.

31. Kazakos GM, Cobb CM, Morrison SL, Baker BF, and Killoy WK, Gingival response to sub-gingival placement of monolithic tetracycline impregnated fibers. Microscopic observation. Int. J. Petriodontics Restorative Dent., 1993, 63, 849.

32. Saxen L, and Asikainen S, Metronidazole in the treatment of localized juvenile periodontitis. K. Clin. Periodontal, 1993, 20, 166.

33. Steenbergh, V and Bery P, Sub-gingival minocycline ointment in moderate to severe chronic adult periodontitis. J. Periodontal, 1993, 64, 637.

34. Stoltze K and Micheal S, Systemic absorption of metronidazole after application of metronidazole dental gel. J. Clin. Periodontal, 1992, 19, 693.

35. Stortze K and Micheal S, Concentration of metronidazole in periodontal pockets after application of metronidazole 25% dental gel. J. Clin Periodontal., 1992, 19, 698.